

VOLUME 277 (2002) PAGES 43623–43630

DOI 10.1074/jbc.AAC120.016177

Correction: Cyclic AMP induces transactivation of the receptors for epidermal growth factor and nerve growth factor, thereby modulating activation of MAP kinase, Akt, and neurite outgrowth in PC12 cells.

Albrecht Piiper, Ivan Dikic, Manfred P. Lutz, Jürgen Leser, Bernd Kronenberger, Robert Elez, Henning Cramer, Werner Müller-Esterl, and Stefan Zeuzem

In Fig. 1B, in the lane marked *EGF*, the result of a similar experiment on the EGF-induced tyrosine phosphorylation of EGFR in the presence of H-89 was shown, which was published elsewhere (Piiper *et al.* (2003) *Biochem. Biophys. Res. Commun.* **301**, 848–854; Fig. 1A, lane *H89+EGF*). The revised Fig. 1B shows the correct lane of the EGF-induced phosphorylation of EGFR in the absence of H-89 that was taken from the same original blot as the 8-Br-cAMP-induced EGFR phosphorylation after 0, 3, and 5 min, respectively. This error does not affect the validity of the results and the conclusions of our study; the authors apologize for any inconvenience this might have caused for the readers.




Figure 1B.

Correction: Cyclic AMP induces transactivation of the receptors for epidermal growth factor and nerve growth factor, thereby modulating activation of MAP kinase, Akt, and neurite outgrowth in PC12 cells.

Albrecht Piiper, Ivan Dikic, Manfred P. Lutz, Jürgen Leser, Bernd Kronenberger, Robert Elez, Henning Cramer, Werner Müller-Esterl and Stefan Zeuzem

J. Biol. Chem. 2020, 295:14792.

doi: 10.1074/jbc.AAC120.016177

Access the most updated version of this article at <http://www.jbc.org/content/295/43/14792>

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 0 references, 0 of which can be accessed free at <http://www.jbc.org/content/295/43/14792.full.html#ref-list-1>